

A FAMILY OF SUMMATION FORMULAS INVOLVING GENERALIZED HARMONIC NUMBERS

¹CHUANAN WEI, ²QINGLUN YAN, ³DIANXUAN GONG

¹*Department of Information Technology
Hainan Medical College, Haikou 571101, China*

²*College of Mathematics and Physics
Nanjing University of Posts and Telecommunications, Nanjing 210046, China*

³*College of Sciences
Hebei Polytechnic University, Tangshan 063009, China*

ABSTRACT. Combining the derivative operator with a binomial sum from the telescoping method, we establish a family of summation formulas involving generalized harmonic numbers.

1. INTRODUCTION

For $x \in \mathbb{C}$ and $l, n \in \mathbb{N}_0$, define the functions $H_n^{(l)}(x)$ by

$$H_0^{(l)}(x) = 0 \quad \text{and} \quad H_n^{(l)}(x) = \sum_{k=1}^n \frac{1}{(x+k)^l} \quad \text{with} \quad n = 1, 2, \dots.$$

Fixing $x = 0$ in the functions just mentioned, we obtain the generalized harmonic numbers:

$$H_0^{(l)} = 0 \quad \text{and} \quad H_n^{(l)} = \sum_{k=1}^n \frac{1}{k^l} \quad \text{with} \quad n = 1, 2, \dots.$$

When $l = 1$, they reduce to the classical harmonic numbers:

$$H_0 = 0 \quad \text{and} \quad H_n = \sum_{k=1}^n \frac{1}{k} \quad \text{with} \quad n = 1, 2, \dots.$$

There exist many elegant identities involving generalized harmonic numbers. They can be found in the papers [1]-[8].

For a differentiable function $f(x)$, define the derivative operator $\mathcal{D}_x$ by

$$\mathcal{D}_x f(x) = \frac{d}{dx} f(x).$$

Then it is not difficult to show the following two derivatives:

$$\begin{aligned} \mathcal{D}_x \binom{x+n}{n} &= \binom{x+n}{n} H_n(x), \\ \mathcal{D}_x H_n^{(l)}(x) &= -l H_n^{(l+1)}(x). \end{aligned}$$

2010 Mathematics Subject Classification: Primary 05A19 and Secondary 40A25

Key words and phrases. Telescoping method; Derivative operator; Generalized harmonic numbers.

Email addresses: weichuanan@yahoo.com.cn (C. Wei), yanqinglun@yahoo.com.cn (Q. Yan), gongdianxuan@yahoo.com.cn (D. Gong).

For a complex sequence $\{\tau_k\}_{k \in \mathbb{Z}}$, define the difference operator by

$$\nabla \tau_k = \tau_k - \tau_{k-1}.$$

Then we have the following relation:

$$\nabla \frac{\binom{y+k+1}{k}}{\binom{x+k}{k}} = \frac{\binom{y+k}{k}}{\binom{x+k}{k}} \frac{y-x+1}{y+1}.$$

Combining the last equation and the telescoping method:

$$\sum_{k=1}^n \nabla \tau_k = \tau_n - \tau_0,$$

we get the simple binomial sum:

$$\sum_{k=1}^n \frac{\binom{y+k}{k}}{\binom{x+k}{k}} = \frac{\binom{y+n+1}{n}}{\binom{x+n}{n}} \frac{y+1}{y-x+1} - \frac{y+1}{y-x+1}. \quad (1)$$

By means of the derivative operator $\mathcal{D}_x$ and the binomial sum (1), we shall explore systematically closed expressions for the family of sums:

$$\sum_{k=1}^n k^i H_k^{(l)}(x) \text{ with } i, l \in \mathbb{N}_0.$$

When $x = p$ with $p \in \mathbb{N}_0$, they give closed expressions for the following sums:

$$\sum_{k=1}^n k^i H_{p+k}^{(l)}.$$

2. SUMMATION FORMULAS

Theorem 1. *For $x \in \mathbb{C}$ and $l \in \mathbb{N}_0$, there holds the summation formula:*

$$\sum_{k=1}^n H_k^{(l+1)}(x) = (x+n+1)H_n^{(l+1)}(x) - H_n^{(l)}(x).$$

Proof. Applying the derivative operator $\mathcal{D}_x$ to (1), we achieve the identity:

$$\sum_{k=1}^n \frac{\binom{y+k}{k}}{\binom{x+k}{k}} H_k(x) = \frac{y+1}{y-x+1} \frac{\binom{y+n+1}{n}}{\binom{x+n}{n}} \left\{ H_n(x) - \frac{1}{y-x+1} \right\} + \frac{y+1}{(y-x+1)^2}. \quad (2)$$

Letting $y = x$ in (2), we attain the case $l = 0$ of Theorem 1:

$$\sum_{k=1}^n H_k(x) = (x+n+1)H_n(x) - n.$$

Suppose that the following identity

$$\sum_{k=1}^n H_k^{(l+1)}(x) = (x+n+1)H_n^{(l+1)}(x) - H_n^{(l)}(x)$$

is true. Applying the derivative operator $\mathcal{D}_x$ to the last equation, we have

$$\sum_{k=1}^n H_k^{(l+2)}(x) = (x+n+1)H_n^{(l+2)}(x) - H_n^{(l+1)}(x).$$

This proves Theorem 1 inductively. □

Making $x = p$ in Theorem 1, we get the following equation.

Corollary 2. For $l, p \in \mathbb{N}_0$, there holds the summation formula:

$$\sum_{k=1}^n H_{p+k}^{\langle l+1 \rangle} = (p+n+1)H_{p+n}^{\langle l+1 \rangle} - (p+1)H_p^{\langle l+1 \rangle} - H_{p+n}^{\langle l \rangle} + H_p^{\langle l \rangle}.$$

3. SUMMATION FORMULAS WITH THE FACTOR k

Setting $y = x + 1$ in (2) and considering the relation:

$$\sum_{k=1}^n \frac{\binom{x+1+k}{k}}{\binom{x+k}{k}} H_k(x) = \sum_{k=1}^n H_k(x) + \frac{1}{x+1} \sum_{k=1}^n k H_k(x),$$

we gain the following equation by using Theorem 1.

Proposition 3. For $x \in \mathbb{C}$, there holds the summation formula:

$$\sum_{k=1}^n k H_k(x) = \frac{(x+n+1)(n-x)}{2} H_n(x) + \frac{(2x-n+1)n}{4}.$$

Corollary 4 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 3).

$$\sum_{k=1}^n k H_{p+k} = \frac{(n-p)(p+n+1)}{2} H_{p+n} + \frac{p(p+1)}{2} H_p - \frac{n(n-2p-1)}{4}.$$

Theorem 5. For $x \in \mathbb{C}$ and $l \in \mathbb{N}_0$, there holds the summation formula:

$$\sum_{k=1}^n k H_k^{\langle l+2 \rangle}(x) = \frac{(x+n+1)(n-x)}{2} H_n^{\langle l+2 \rangle}(x) + \frac{2x+1}{2} H_n^{\langle l+1 \rangle}(x) - \frac{H_n^{\langle l \rangle}(x)}{2}.$$

Proof. Applying the derivative operator $\mathcal{D}_x$ to Proposition 3, we achieve the case $l = 0$ of Theorem 5:

$$\sum_{k=1}^n k H_k^{\langle 2 \rangle}(x) = \frac{(x+n+1)(n-x)}{2} H_n^{\langle 2 \rangle}(x) + \frac{2x+1}{2} H_n(x) - \frac{n}{2}.$$

Suppose that the following identity

$$\sum_{k=1}^n k H_k^{\langle l+2 \rangle}(x) = \frac{(x+n+1)(n-x)}{2} H_n^{\langle l+2 \rangle}(x) + \frac{2x+1}{2} H_n^{\langle l+1 \rangle}(x) - \frac{H_n^{\langle l \rangle}(x)}{2}.$$

is true. Applying the derivative operator $\mathcal{D}_x$ to the last equation, we have

$$\sum_{k=1}^n k H_k^{\langle l+3 \rangle}(x) = \frac{(x+n+1)(n-x)}{2} H_n^{\langle l+3 \rangle}(x) + \frac{2x+1}{2} H_n^{\langle l+2 \rangle}(x) - \frac{H_n^{\langle l+1 \rangle}(x)}{2}.$$

This proves Theorem 5 inductively. $\square$

Taking $x = p$ in Theorem 5, we attain the following equation.

Corollary 6. For $l, p \in \mathbb{N}_0$, there holds the summation formula:

$$\begin{aligned} \sum_{k=1}^n k H_{p+k}^{\langle l+2 \rangle} &= \frac{(p+n+1)(n-p)}{2} H_{p+n}^{\langle l+2 \rangle} + \frac{p(p+1)}{2} H_p^{\langle l+2 \rangle} \\ &\quad + \frac{2p+1}{2} (H_{p+n}^{\langle l+1 \rangle} - H_p^{\langle l+1 \rangle}) - \frac{H_{p+n}^{\langle l \rangle} - H_p^{\langle l \rangle}}{2}. \end{aligned}$$

4. SUMMATION FORMULAS WITH THE FACTOR k^2

Letting $y = x + 2$ in (2) and considering the relation:

$$\begin{aligned} \sum_{k=1}^n \frac{\binom{x+2+k}{k}}{\binom{x+k}{k}} H_k(x) &= \sum_{k=1}^n H_k(x) + \frac{2x+3}{(x+1)(x+2)} \sum_{k=1}^n k H_k(x) \\ &\quad + \frac{1}{(x+1)(x+2)} \sum_{k=1}^n k^2 H_k(x), \end{aligned}$$

we get the following equation by using Theorem 1 and Proposition 3.

Proposition 7. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^2 H_k(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n(x) \\ &\quad - \frac{(12x^2 + 12x - 6xn + 4n^2 - 3n - 1)n}{36}. \end{aligned}$$

Corollary 8 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 7).

$$\begin{aligned} \sum_{k=1}^n k^2 H_{p+k} &= \frac{(p+n+1)(2n^2 + n - 2pn + p + 2p^2)}{6} H_{p+n} \\ &\quad - \frac{p(p+1)(2p+1)}{6} H_p - \frac{n(4n^2 - 3n - 6pn + 12p + 12p^2 - 1)}{36}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 7, we gain the following equation.

Proposition 9. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^2 H_k^{(2)}(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n^{(2)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{6} H_n(x) + \frac{(4x + 2 - n)n}{6}. \end{aligned}$$

Corollary 10 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 9).

$$\begin{aligned} \sum_{k=1}^n k^2 H_{p+k}^{(2)} &= \frac{p(p+1)(2p+1) + n(n+1)(2n+1)}{6} H_{p+n}^{(2)} - \frac{p(p+1)(2p+1)}{6} H_p^{(2)} \\ &\quad - \frac{6p^2 + 6p + 1}{6} (H_{p+n} - H_p) + \frac{(4p + 2 - n)n}{6}. \end{aligned}$$

Theorem 11. *For $x \in \mathbb{C}$ and $l \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^2 H_k^{(l+3)}(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n^{(l+3)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{6} H_n^{(l+2)}(x) + \frac{2x+1}{2} H_n^{(l+1)}(x) - \frac{H_n^{(l)}(x)}{3}. \end{aligned}$$

Proof. Applying the derivative operator $\mathcal{D}_x$ to Proposition 9, we achieve the case $l = 0$ of Theorem 11:

$$\begin{aligned} \sum_{k=1}^n k^2 H_k^{(3)}(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n^{(3)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{6} H_n^{(2)}(x) + \frac{2x+1}{2} H_n(x) - \frac{n}{3}. \end{aligned}$$

Suppose that the following identity

$$\begin{aligned} \sum_{k=1}^n k^2 H_k^{(l+3)}(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n^{(l+3)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{6} H_n^{(l+2)}(x) + \frac{2x+1}{2} H_n^{(l+1)}(x) - \frac{H_n^{(l)}(x)}{3} \end{aligned}$$

is true. Applying the derivative operator $\mathcal{D}_x$ to the last equation, we have

$$\begin{aligned} \sum_{k=1}^n k^2 H_k^{(l+4)}(x) &= \frac{x(x+1)(2x+1) + n(n+1)(2n+1)}{6} H_n^{(l+4)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{6} H_n^{(l+3)}(x) + \frac{2x+1}{2} H_n^{(l+2)}(x) - \frac{H_n^{(l+1)}(x)}{3}. \end{aligned}$$

This proves Theorem 11 inductively. $\square$

Making $x = p$ in Theorem 11, we attain the following equation.

Corollary 12. *For $l, p \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^2 H_{p+k}^{(l+3)} &= \frac{p(p+1)(2p+1) + n(n+1)(2n+1)}{6} H_{p+n}^{(l+3)} \\ &\quad - \frac{p(p+1)(2p+1)}{6} H_p^{(l+3)} - \frac{6p^2 + 6p + 1}{6} (H_{p+n}^{(l+2)} - H_p^{(l+2)}) \\ &\quad + \frac{2p+1}{2} (H_{p+n}^{(l+1)} - H_p^{(l+1)}) - \frac{H_{p+n}^{(l)} - H_p^{(l)}}{3}. \end{aligned}$$

5. SUMMATION FORMULAS WITH THE FACTOR k^3

Setting $y = x + 3$ in (2) and considering the relation:

$$\begin{aligned} \sum_{k=1}^n \frac{\binom{x+3+k}{k}}{\binom{x+k}{k}} H_k(x) &= \sum_{k=1}^n H_k(x) + \frac{3x^2 + 12x + 11}{(x+1)(x+2)(x+3)} \sum_{k=1}^n k H_k(x) \\ &\quad + \frac{3}{(x+1)(x+3)} \sum_{k=1}^n k^2 H_k(x) \\ &\quad + \frac{1}{(x+1)(x+2)(x+3)} \sum_{k=1}^n k^3 H_k(x), \end{aligned}$$

we get the following equation by using Theorem 1, Proposition 3 and Proposition 7.

Proposition 13. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^3 H_k(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n(x) \\ &\quad - \frac{(12x^3 + 18x^2 - 6nx^2 + 2x - 6xn + 4n^2x - 2 + 3n + 2n^2 - 3n^3)n}{48}. \end{aligned}$$

Corollary 14 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 13).

$$\begin{aligned} \sum_{k=1}^n k^3 H_{p+k} &= \frac{(n-p)(p+n+1)(p^2+p+n+n^2)}{4} H_{p+n} + \frac{p^2(p+1)^2}{4} H_p \\ &\quad - \frac{(12p^3 + 18p^2 - 6np^2 + 2p - 6pn + 4n^2p - 2 + 3n + 2n^2 - 3n^3)n}{48}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 13, we gain the following equation.

Proposition 15. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(2)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(2)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n(x) - \frac{(18x^2+18x-6nx+1-3n+2n^2)n}{24}. \end{aligned}$$

Corollary 16 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 15).

$$\begin{aligned} \sum_{k=1}^n k^3 H_{p+k}^{(2)} &= \frac{(n-p)(p+n+1)(p^2+p+n+n^2)}{4} H_{p+n}^{(2)} + \frac{p^2(p+1)^2}{4} H_p^{(2)} \\ &+ \frac{p(p+1)(2p+1)}{2} (H_{p+n} - H_p) - \frac{(18p^2+18p-6np+1-3n+2n^2)n}{24}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 15, we achieve the following equation.

Proposition 17. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(3)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(3)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n^{(2)}(x) - \frac{6x^2+6x+1}{4} H_n(x) + \frac{(6x+3-n)n}{8}. \end{aligned}$$

Corollary 18 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 17).

$$\begin{aligned} \sum_{k=1}^n k^3 H_{p+k}^{(3)} &= \frac{(n-p)(p+n+1)(p^2+p+n+n^2)}{4} H_{p+n}^{(3)} + \frac{p^2(p+1)^2}{4} H_p^{(3)} \\ &+ \frac{p(p+1)(2p+1)}{2} (H_{p+n}^{(2)} - H_p^{(2)}) - \frac{6p^2+6p+1}{4} (H_{p+n} - H_p) \\ &+ \frac{(6p+3-n)n}{8}. \end{aligned}$$

Theorem 19. *For $x \in \mathbb{C}$ and $l \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(l+4)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(l+4)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n^{(l+3)}(x) - \frac{6x^2+6x+1}{4} H_n^{(l+2)}(x) \\ &+ \frac{2x+1}{2} H_n^{(l+1)}(x) - \frac{H_n^{(l)}(x)}{4}. \end{aligned}$$

Proof. Applying the derivative operator $\mathcal{D}_x$ to Proposition 17, we attain the case $l = 0$ of Theorem 19:

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(4)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(4)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n^{(3)}(x) - \frac{6x^2+6x+1}{4} H_n^{(2)}(x) \\ &+ \frac{2x+1}{2} H_n(x) - \frac{n}{4}. \end{aligned}$$

Suppose that the following identity

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(l+4)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(l+4)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n^{(l+3)}(x) - \frac{6x^2+6x+1}{4} H_n^{(l+2)}(x) \\ &+ \frac{2x+1}{2} H_n^{(l+1)}(x) - \frac{H_n^{(l)}(x)}{4} \end{aligned}$$

is true. Applying the derivative operator $\mathcal{D}_x$ to the last equation, we have

$$\begin{aligned} \sum_{k=1}^n k^3 H_k^{(l+5)}(x) &= \frac{(n-x)(x+n+1)(x^2+x+n+n^2)}{4} H_n^{(l+5)}(x) \\ &+ \frac{x(x+1)(2x+1)}{2} H_n^{(l+4)}(x) - \frac{6x^2+6x+1}{4} H_n^{(l+3)}(x) \\ &+ \frac{2x+1}{2} H_n^{(l+2)}(x) - \frac{H_n^{(l+1)}(x)}{4}. \end{aligned}$$

This proves Theorem 19 inductively. $\square$

Taking $x = p$ in Theorem 19, we get the following equation.

Corollary 20. *For $l, p \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^3 H_{p+k}^{(l+4)} &= \frac{(n-p)(p+n+1)(p^2+p+n+n^2)}{4} H_{p+n}^{(l+4)} + \frac{p^2(p+1)^2}{4} H_p^{(l+4)} \\ &+ \frac{p(p+1)(2p+1)}{2} (H_{p+n}^{(l+3)} - H_p^{(l+3)}) - \frac{6p^2+6p+1}{4} (H_{p+n}^{(l+2)} - H_p^{(l+2)}) \\ &+ \frac{2p+1}{2} (H_{p+n}^{(l+1)} - H_p^{(l+1)}) - \frac{H_{p+n}^{(l)} - H_p^{(l)}}{4}. \end{aligned}$$

6. SUMMATION FORMULAS WITH THE FACTOR k^4

Letting $y = x + 4$ in (2) and considering the relation:

$$\begin{aligned} \sum_{k=1}^n \frac{\binom{x+4+k}{k}}{\binom{x+k}{k}} H_k(x) &= \sum_{k=1}^n H_k(x) + \frac{2(2x+5)(x^2+5x+5)}{(x+1)(x+2)(x+3)(x+4)} \sum_{k=1}^n k H_k(x) \\ &+ \frac{6x^2+30x+35}{(x+1)(x+2)(x+3)(x+4)} \sum_{k=1}^n k^2 H_k(x) \\ &+ \frac{4x+10}{(x+1)(x+2)(x+3)(x+4)} \sum_{k=1}^n k^3 H_k(x) \\ &+ \frac{1}{(x+1)(x+2)(x+3)(x+4)} \sum_{k=1}^n k^4 H_k(x), \end{aligned}$$

we gain the following equation by using Theorem 1, Proposition 3, Proposition 7 and Proposition 13.

Proposition 21. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_k(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n(x) \\ &\quad - \frac{(72n^4 - 45n^3 - 130n^2 + 75n + 28)n}{1800} \\ &\quad - \frac{(12x^3 + 24x^2 + 7x - 6nx^2 - 9nx + 4n^2x - 5 + 2n + 4n^2 - 3n^3)nx}{60}. \end{aligned}$$

Corollary 22 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 21).

$$\begin{aligned} \sum_{k=1}^n k^4 H_{p+k} &= \frac{6p^5 + 15p^4 + 10p^3 - p - n + 10n^3 + 15n^4 + 6n^5}{30} H_{p+n} \\ &\quad - \frac{6p^5 + 15p^4 + 10p^3 - p}{30} H_p - \frac{(72n^4 - 45n^3 - 130n^2 + 75n + 28)n}{1800} \\ &\quad - \frac{(12p^3 + 24p^2 + 7p - 6np^2 - 9np + 4n^2p - 5 + 2n + 4n^2 - 3n^3)np}{60}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 21, we achieve the following equation.

Proposition 23. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{(2)}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{(2)}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n(x) \\ &\quad + \frac{(48x^3 + 72x^2 - 18nx^2 + 14x - 18nx + 8n^2x - 5 + 2n + 4n^2 - 3n^3)n}{60}. \end{aligned}$$

Corollary 24 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 23).

$$\begin{aligned} \sum_{k=1}^n k^4 H_{p+k}^{(2)} &= \frac{6p^5 + 15p^4 + 10p^3 - p - n + 10n^3 + 15n^4 + 6n^5}{30} H_{p+n}^{(2)} \\ &\quad - \frac{6p^5 + 15p^4 + 10p^3 - p}{30} H_p^{(2)} - \frac{30p^2(p+1)^2 - 1}{30} (H_{p+n} - H_p) \\ &\quad + \frac{(48p^3 + 72p^2 - 18np^2 + 14p - 18np + 8n^2p - 5 + 2n + 4n^2 - 3n^3)n}{60}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 23, we attain the following equation.

Proposition 25. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{(3)}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{(3)}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{(2)}(x) + x(x+1)(2x+1)H_n(x) \\ &\quad - \frac{(72x^2 + 72x - 18nx + 7 - 9n + 4n^2)n}{60}. \end{aligned}$$

Corollary 26 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 25).

$$\begin{aligned} \sum_{k=1}^n k^4 H_{p+k}^{(3)} &= \frac{6p^5 + 15p^4 + 10p^3 - p - n + 10n^3 + 15n^4 + 6n^5}{30} H_{p+n}^{(3)} \\ &\quad - \frac{6p^5 + 15p^4 + 10p^3 - p}{30} H_p^{(3)} - \frac{30p^2(p+1)^2 - 1}{30} (H_{p+n}^{(2)} - H_p^{(2)}) \\ &\quad + p(p+1)(2p+1)(H_{p+n} - H_p) - \frac{(72p^2 + 72p - 18np + 7 - 9n + 4n^2)n}{60}. \end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Proposition 25, we get the following equation.

Proposition 27. *For $x \in \mathbb{C}$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{(4)}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{(4)}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{(3)}(x) + x(x+1)(2x+1)H_n^{(2)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{3} H_n(x) + \frac{(8x + 4 - n)n}{10}. \end{aligned}$$

Corollary 28 ($x = p$ with $p \in \mathbb{N}_0$ in Proposition 27).

$$\begin{aligned} \sum_{k=1}^n k^4 H_{p+k}^{(4)} &= \frac{6p^5 + 15p^4 + 10p^3 - p - n + 10n^3 + 15n^4 + 6n^5}{30} H_{p+n}^{(4)} \\ &\quad - \frac{6p^5 + 15p^4 + 10p^3 - p}{30} H_p^{(4)} - \frac{30p^2(p+1)^2 - 1}{30} (H_{p+n}^{(3)} - H_p^{(3)}) \\ &\quad + p(p+1)(2p+1)(H_{p+n}^{(2)} - H_p^{(2)}) - \frac{6p^2 + 6p + 1}{3} (H_{p+n} - H_p) \\ &\quad + \frac{(8p + 4 - n)n}{10}. \end{aligned}$$

Theorem 29. *For $x \in \mathbb{C}$ and $l \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{(l+5)}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{(l+5)}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{(l+4)}(x) + x(x+1)(2x+1)H_n^{(l+3)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{3} H_n^{(l+2)}(x) + \frac{2x+1}{2} H_n^{(l+1)}(x) - \frac{H_n^{(l)}(x)}{5}. \end{aligned}$$

Proof. Applying the derivative operator $\mathcal{D}_x$ to Proposition 27, we gain the case $l = 0$ of Theorem 29:

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{(5)}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{(5)}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{(4)}(x) + x(x+1)(2x+1)H_n^{(3)}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{3} H_n^{(2)}(x) + \frac{2x+1}{2} H_n(x) - \frac{n}{5}. \end{aligned}$$

Suppose that the identity

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{\langle l+5 \rangle}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{\langle l+5 \rangle}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{\langle l+4 \rangle}(x) + x(x+1)(2x+1) H_n^{\langle l+3 \rangle}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{3} H_n^{\langle l+2 \rangle}(x) + \frac{2x+1}{2} H_n^{\langle l+1 \rangle}(x) - \frac{H_n^{\langle l \rangle}(x)}{5} \end{aligned}$$

is true. Applying the derivative operator $\mathcal{D}_x$ to the last equation, we have

$$\begin{aligned} \sum_{k=1}^n k^4 H_k^{\langle l+6 \rangle}(x) &= \frac{6x^5 + 15x^4 + 10x^3 - x - n + 10n^3 + 15n^4 + 6n^5}{30} H_n^{\langle l+6 \rangle}(x) \\ &\quad - \frac{30x^2(x+1)^2 - 1}{30} H_n^{\langle l+5 \rangle}(x) + x(x+1)(2x+1) H_n^{\langle l+4 \rangle}(x) \\ &\quad - \frac{6x^2 + 6x + 1}{3} H_n^{\langle l+3 \rangle}(x) + \frac{2x+1}{2} H_n^{\langle l+2 \rangle}(x) - \frac{H_n^{\langle l+1 \rangle}(x)}{5}. \end{aligned}$$

This proves Theorem 29 inductively. $\square$

Making $x = p$ in Theorem 29, we obtain the following equation.

Corollary 30. *For $l, p \in \mathbb{N}_0$, there holds the summation formula:*

$$\begin{aligned} \sum_{k=1}^n k^4 H_{p+k}^{\langle l+5 \rangle} &= \frac{6p^5 + 15p^4 + 10p^3 - p - n + 10n^3 + 15n^4 + 6n^5}{30} H_{p+n}^{\langle l+5 \rangle} \\ &\quad - \frac{6p^5 + 15p^4 + 10p^3 - p}{30} H_{p+n}^{\langle l+5 \rangle} - \frac{30p^2(p+1)^2 - 1}{30} (H_{p+n}^{\langle l+4 \rangle} - H_p^{\langle l+4 \rangle}) \\ &\quad + p(p+1)(2p+1) (H_{p+n}^{\langle l+3 \rangle} - H_p^{\langle l+3 \rangle}) - \frac{6p^2 + 6p + 1}{3} (H_{p+n}^{\langle l+2 \rangle} - H_p^{\langle l+2 \rangle}) \\ &\quad + \frac{2p+1}{2} (H_{p+n}^{\langle l+1 \rangle} - H_p^{\langle l+1 \rangle}) - \frac{H_{p+n}^{\langle l \rangle} - H_p^{\langle l \rangle}}{5}. \end{aligned}$$

Remark: Further summation formulas with the factor k^i , where i is a positive integer greater than 4, can also be derived in the same way. Considering that the resulting identities will become more complicated, we shall not lay out them here.

REFERENCES

- [1] G. E. Andrews, K. Uchimura, Identities in combinatorics IV: differentiation and harmonic numbers, *Utilitas Math.* 28 (1985) 265-269.
- [2] Y. Chen, Q. Hou, H. Jin, The Abel-Zeilberge algorithm, *Electron. J. Comb.* 18 (2011) #P17.
- [3] F. Chyzak, An extension of Zeilberger's fast algorithm to general holonomic functions, *Discrete Math.* 217 (2000) 115-134.
- [4] H. Liu, W. Wang, Harmonic number identities via hypergeometric series and Bell polynomials, *Integral Transforms Spec. Funct.* 23 (2012) 49-68.
- [5] P. Paule, C. Schneider, Computer proofs of a new family of harmonic number identities, *Adv. Appl. Math.* 31 (2003) 359-378.
- [6] C. Schneider, Symbolic summation assists Combinatorics, *Sém. Lothar. Combin.* 56 (2006) Article B56b.
- [7] A. Sofu, Some more identities involving rational sums, *Appl. Anal. Discr. Math.* 2 (2008) 56-66.
- [8] D. Zheng, Further summation formulae related to generalized harmonic numbers, *J. Math. Anal. Appl.* 335 (2007) 692-706.